

**Marcin J. Małuszyński\***

## **THALLIUM IN ENVIRONMENT**

### **TAL W ŚRODOWISKU**

**Słowa kluczowe:** tal, środowisko, zanieczyszczenie środowiska.

**Key words:** thallium, environment, environmental pollution.

*Tal w środowiskach geochemicznych występuje na trzech stopniach utlenienia choć najczęściej jako  $Tl^+$ . Jest on silnie sorbowany przez minerały ilaste, wodorotlenki Fe i Mn, oraz substancję organiczną, co wskazuje na potencjalną zdolność tego pierwiastka do przechodzenia ze środowiska glebowego do łańcucha troficznego.*

*Tal uznawany jest jako pierwiastek toksyczny, którego oddziaływanie na organizmy kręgowców może być zdaniem specjalistów bardziej niebezpieczne niż rtęci, kadmu czy ołowiu. Jest szczególnie łatwo pobierany przez rośliny, a jego szkodliwy wpływ objawia się zahamowaniem rozwoju systemu korzeniowego, karłowaceniem roślin oraz chlorozą. Szkodliwość talu objawia się także zahamowaniem zawiązywania nasion, oraz zmniejszeniem zdolności kiełkowania. Tal jest pierwiastkiem toksycznym także dla organizmów zwierząt i ludzi. Jest on równie toksyczny przy pobieraniu drogą pokarmową, jak i wskutek wdychania pyłów lub par oraz wchłaniania przez skórę. Charakterystycznym objawem zatrucia talu jest łysienie. W organizmach zatrutych występują zmiany psychiczne oraz uszkodzenia układu naczyniowo-sercowego.*

#### **1. INTRODUCTION**

Thallium atomic number 81, was discovered spectroscopically in 1861 by Crookes. The element was named after the green spectral line, which identified the element (Greek „thallos”, green twig). The metal was isolated both by Crookes and Lamy in 1862. They had been

---

\* *Dr inż. Marcin J. Małuszyński – Katedra Inżynierii Wodnej i Rekultywacji Środowiska, Wydział Inżynierii i Kształtowania Środowiska, Szkoła Główna Gospodarstwa Wiejskiego, ul. Nowoursynowska 166, 02-787 Warszawa, tel.: 22 593 53 39; e-mail: marcin\_maluszynski@sggw.pl*

expecting to isolate tellurium after removing selenium from the byproducts from a commercial sulphuric acid factory but instead found the new element thallium [Winter 2007].

Thallium occurs mainly in association with potassium minerals, such as sylvite and polucite, in clays, soils and granites. Thallium minerals are well known too; they are rare, but a few are known, such as crookesite, lorandite, christite, avicennite, ellisite or sicherite. They contain 16–60% of thallium, namely as sulphides or selenides in complexes with antimony, arsenic, copper, lead and silver [Anderson et al. 1999; Kazantzis 2000; Xiao et al. 2004a].

Thallium is also produced as a byproduct from zinc and lead refining plants, as well as from particular sulphuric acid factories [Winter 2007].

The natural release of thallium to environment occurs primarily as a result of weathering of rocks and soil erosion. During weathering, Tl is readily mobilized and transported together with alkaline metals. However, Tl is most often sorbed by clays, Fe/Mn oxides and organic matter. The largest anthropogenic source of Tl is coal combustion, but some amount of this element may be released by heavy metal smelting, cement industry and refining processes [Kabata-Pendias, Pendias 2001].

Thallium chemical behaviour resembles the heavy metals (Pb, Au and Ag) on the one hand and the alkali metals (K, Rb, Cs) on the other [Rao et al. 2008].

## 2. THALLIUM IN THE AIR

Some anthropogenic activities, such as cement factories using iron oxides with high Tl contents, coal combustion and ferrous and non-ferrous metallurgy, can release Tl into the biosphere and may increase its availability [Tremel et al. 1997b, Hršak et al. 2003].

Determination of thallium in Zagreb air was performed by Hršak et al. [2003]. The concentration of thallium in 1998 ranged between 0 to 18.78 ng·m<sup>-3</sup> in city centre while in residential area it was lower (0–5.04 ng·m<sup>-3</sup>). In the year 2000 same researchers found lower Tl concentration in city centre (0–2.17 ng·m<sup>-3</sup>) than in residential area (0–8.90 ng·m<sup>-3</sup>).

## 3. THALLIUM IN SOILS

Thallium is found in alloys, minerals, biological samples, and geological materials. Its content of soils depends largely on the geological origin of the parent material. In general, thallium concentrations in surface soils range from 0.1 to 2 µg·g<sup>-1</sup>, with most reported concentrations being <1 µg·g<sup>-1</sup>. Its concentration in arable soils is between 0.29 and 1.54 µg·g<sup>-1</sup>. High concentrations of the metal (up to 55 µg·g<sup>-1</sup>) are common in soils derived from limestone, marble, or granite, because these minerals contain appreciable amounts of sulfur, which in turn has high affinity for Tl. Thallium salts are highly water soluble, but can be immobilized by vermiculite once incorporated into soil [Martin, Kaplan 1998, Tremel et al. 1997b, Qi et al. 1992, Jacobson et al. 2005].

Concentration of thallium in soils of the EuroRegion Neisse was analysed by Heim et al. [2002]. The median concentration in this area was  $0.5 \text{ mg}\cdot\text{kg}^{-1}$ . This compares to a median  $0.3 \text{ mg}\cdot\text{kg}^{-1}$  TI in French topsoils reported by Tremel et al. [1997a] and to  $0.44 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  in soils from the south-eastern part of the Silesian-Cracovian zinc-lead ore deposit region [Jakubowska et al. 2007]. Soil samples collected from grasslands in Czech Republic by Vaněk et al. [2009] contained about  $1.1 \text{ mg}\cdot\text{kg}^{-1}$  of thallium. Thallium concentration in soils from region of Kosovska Mitrovica varied between  $0.10$  and  $5.06 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  [Borgna et al. 2009]. In Scania, the most southern province of Sweden, concentration of thallium in soils was about  $0.32 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  average [Tyler 2004b]. The results of Qi et al. [1992] research show that soils from 34 provinces of China contains  $0.292$ – $1.172 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ . They reported that soils developed on limestones have the maximum concentration of thallium. They also observed negative correlation between TI content and pH of the soils and positive one between TI content and organic matter of soils.

Yang et al. [2005] also analysed the concentration of the thallium in the Chinese soils. The concentration of TI in the background soil ranging from  $1.63$  to  $2.02 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ , while in the upper layer of soils which are spatially closed to the slag pile high concentration of thallium (up to  $15.4 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ ) was observed.

#### 4. THALLIUM IN WATERS

The solubility of thallos compounds is relatively high so that monovalent thallium is readily transported through aqueous routes into the environment. Thallium is included in the USEPA list of priority toxic pollutants and fixed  $2 \text{ }\mu\text{g}\cdot\text{L}^{-1}$  [EPA] as maximum contaminant level in drinking water.

Peter and Viraraghavan [2005] present that drinking water sometimes may contain  $7.2 \text{ }\mu\text{g TI}\cdot\text{L}^{-1}$ , while concentration of thallium in groundwater ranged from  $20$  to  $24 \text{ }\mu\text{g}\cdot\text{L}^{-1}$ .

Results of Xiao et al. [2004b] research indicates that TI concentration in deep groundwater is higher ( $13$ – $1100 \text{ }\mu\text{g}\cdot\text{L}^{-1}$ ) and its decreasing with distance away from mineralized area to background levels ( $<0.005 \text{ }\mu\text{g}\cdot\text{L}^{-1}$ ).

Jakubowska et al. [2007] reported that water samples collected from Biała Przemsza River flowing through the zinc-lead ore exploration area contained  $2.07 \text{ }\mu\text{g}\cdot\text{L}^{-1}$  TI. Water samples collected by Krasnodębska-Ostręga et al. [2005] from another Polish zinc-lead ore exploration and smelting area (near Bukowno) contained lower amounts of thallium ( $0.02$ – $0.15 \text{ }\mu\text{g}\cdot\text{L}^{-1}$ ).

#### 5. THALLIUM IN PLANTS

Thallium is easily available to plants, and when concentrated in roots up to  $2 \text{ mg}\cdot\text{kg}^{-1}$ , can inhibit the germination, plant growth, and chlorophyll content. Plants especially sensitive to in-

creased levels of thallium are the Leguminosae species, cereals, tobacco, and buckwheat. Microorganisms are reported to be relatively sensitive to thallium, and therefore the inhibition of nitrate formation in Tl-polluted soils may have an agronomic impact [Rao et al. 2008].

As far as relatively high content of Tl is appearing in soils, a potential risk for humans can arise at levels around  $1 \text{ mg}\cdot\text{kg}^{-1}$ . Especially growing of certain crops e.g. *Brassicaceae* can pose a risk for the food chains because of elevated accumulation of Tl in plant tissues [Pavlickova et al. 2005].

Uptake by plants can lead to substantial accumulation of this heavy metal. Tremel et al. [1997b] determined the content of the thallium in plants. Both grains wheat and maize contained less than  $4 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  of thallium. Maize stalk had quite a high Tl concentration ( $343 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ ) and elevated Tl concentrations ( $20 \text{ mg}\cdot\text{kg}^{-1}$ ) they found in shoots of rape.

Thallium exhibits the surprising ability to be preconcentrated by green cabbage which significantly increases the daily uptake of thallium by inhabitants of the polluted area [Xiao et al. 2004a]. Roots of some plants also exhibit a high preconcentration of thallium. Tyler [2004a] reports that in Scania (Sweden) roots of *Fagus sylvatica* contain  $0.29 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  of this element, while its concentration in soil was  $0.06 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ . Similar relation was observed by Wierzbicka et al. [2004]. They found that on the waste heap in Bolesław (Poland), on three stands *Plantago lanceolata* accumulated the largest amount of thallium in its roots, on average  $154 (\pm 112) \text{ }\mu\text{g}\cdot\text{g}^{-1}$  (maximum  $321 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ ). The shoots of these plants also contained high thallium levels, on average  $54 (\pm 53) \text{ }\mu\text{g}\cdot\text{g}^{-1}$  (maximum  $180 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ ) Thallium concentrations in soils in all of the tested sites were relatively high and averaged  $43 (\pm 12) \text{ }\mu\text{g}\cdot\text{g}^{-1}$ .

During the research in the EuroRegion Neisse, Heim et al. [2002] found  $0.04\text{--}0.13 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  of thallium in *Pleurozium schreberi*. Similar content was found by Rühling and Tyler [2004] in *Pleurozium schreberi* sampled in 1975 ( $0.15 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ ) and 2000 ( $0.066 \text{ }\mu\text{g}\cdot\text{g}^{-1}$ ) in southern Sweden. They tried to find possible causes of this change, and they conclude that „reductions in anthropogenic dust emissions during recent decades have decreased the atmospheric deposition over northern Europe of this element”.

Anderson et al. [1999] discovered unusually high hyperaccumulation of Tl by the brassicaceae *Iberis intermedia* and *Biscutella laevigata* L. growing over lead-zinc mine tailings at Les Malines near Montpellier in southern France. Both *Biscutella laevigata* and *Iberis intermedia* from Les Malines contained very high levels of Tl of up to 1.4% dry weight in *Biscutella* and 0.4% in *Iberis* in the dry matter, therefore they suggested to use these two species of plants for removing (phytomining) the thallium from the environment.

## 6. THALLIUM IN ORGANISMS OF ANIMALS AND PEOPLE

Thallium is being readily transferred to the human body, from contaminated soils, through food chain, and poses significant threat to the inhabitants of the polluted area [Xiao et al. 2004a].

Most thallium salts are rapidly absorbed through the respiratory and gastrointestinal tracts and through the skin. In humans and animals, thallium circulates in intra- and extra-cellular fluids as a monovalent cation. Its toxicity results from thallium ion mimicking potassium ions in many metabolic processes. Ionic radius of  $Tl^+$  (1.49 Å) is similar to that of the hard cation, potassium  $K^+$  (1.33 Å), so that nondiscriminatory uptake of  $Tl^+$  over  $K^+$  has been suggested as a mechanism of its toxicity to biota. Thallium may also bind with sulfhydryl groups of proteins to inactivate many enzymatic reactions. Thallium causes gastrointestinal irritation and nerve damage when people are exposed to it for relatively short period of time. For long term, thallium has the potential to cause effects such as change in blood chemistry, damage to liver, kidney, intestinal and testicular tissue. In humans, one of the main symptoms of thallium poisoning is rapid (2–3 weeks) and often complete hair loss. Acute poisoning can lead to paralysis, coma, and death. Thallium also has mutagenic, carcinogenic and teratogenic effects [Galván-Arzate and Santamaría 1998, Leonard and Gerber 1997, Moore et al. 1993, Tremel et al. 1997b, Wang et al. 2007].

According to Rusyniak et al. [2002] „thallium has qualities of a perfect criminal poison. Its salts are tasteless, odorless, dissolve completely in liquids, are rapidly absorbed, and defy detection on routine toxicologic screens”. Authors suggesting „although thallium poisoning is an uncommon method of criminal poisoning, it should be considered in any patient with rapidly progressive, unexplained peripheral neuropathy with or without alopecia”.

In order to establish reference values of thallium in tissues of healthy inhabitants living in Central Italy, Sabbioni et al. [1994] determined this element in blood and urine. Mean measured values of thallium in blood and urine were  $0.063 \mu\text{g}\cdot\text{L}^{-1}$  and  $0.066 \mu\text{g}\cdot\text{L}^{-1}$  respectively.

According to Moore et al. [1993] the normal total blood thallium concentration is under  $2 \mu\text{g}\cdot\text{L}^{-1}$  and concentration greater than  $100 \mu\text{g}\cdot\text{L}^{-1}$  are toxic. Urine thallium concentration greater than  $200 \mu\text{g}\cdot\text{L}^{-1}$  are also toxic.

Results reported by several researchers suggesting appearing of the trend in the concentration of thallium in the human body. Concentration of thallium is in rising order: brain ( $0.42\text{--}1.5 \text{ ng}\cdot\text{g}^{-1}$ ) < liver ( $1.5 \text{ ng}\cdot\text{g}^{-1}$ ) < kidney ( $6.1 \text{ ng}\cdot\text{g}^{-1}$ ) < hair ( $7\text{--}650 \text{ ng}\cdot\text{g}^{-1}$ )  $\approx$  bone ( $<0.6 \mu\text{g}\cdot\text{g}^{-1}$ ) < nail ( $1.2 \mu\text{g}\cdot\text{g}^{-1}$ ). This indicates that Tl accumulates preferentially in peripheral organs, for example nails [Engström et al. 2004, Das et al. 2006].

Typical concentrations in animal muscle tissue are: fish ( $0.74\text{--}110.5 \text{ ng}\cdot\text{g}^{-1}$ ), rabbit ( $0.84 \text{ ng}\cdot\text{g}^{-1}$ ), pig ( $1.7 \text{ ng}\cdot\text{g}^{-1}$ ), cattle ( $0.74 \text{ ng}\cdot\text{g}^{-1}$ ) [Engström et al. 2004, Das et al. 2006].

There is little information about Tl contamination in wildlife. Mochizuki et al. [2005] were examined thallium content in wild ducks kidneys and livers. The Tl content ranged from 0.42 to 119.61 and 0.10 to  $33.94 \mu\text{g}\cdot\text{g}^{-1}$  dry weight, respectively. Dmowski et al. [1998] measured high thallium levels in kidneys and livers of small mammals inhabiting the zinc smelter Boleślaw (Southern Poland). The maximum Tl content in kidneys was  $44.06 \mu\text{g}\cdot\text{g}^{-1}$  dry weight, while in livers maximum was  $15.21 \mu\text{g}\cdot\text{g}^{-1}$  dry weight.

## 7. CONCLUSIONS

Thallium is extremely toxic to humans, plants, and animals, and plays no role in their metabolisms. With respect to degree of toxicity thallium ranks alongside Pb, Hg, Cd. The high toxicity of thallium and its compounds made their accurate determination is demanded.

Much further research is required to investigate the possible adverse health effects of thallium following population exposure.

Thallium accumulation demonstrated in plants, fruits and vegetables, that can make big problem because of absence of threshold limits for thallium in soils, agricultural products, feedstuffs and foodstuffs in most countries.

***Acknowledgements: The Preparation and presentation of this article were possible thanks to the financial support of the Ministry of Science and Higher Education (grant no 2P06S01530).***

## REFERENCES

- ANDERSON C.W.N., BROOKS R.R., CHIRUCCI A., LACOSTE C.J., LEBLANS M., ROBINSON B.H., SIMCOCK R., STEWART R.B. 1999. Phytomining for nickel, thallium and gold. *J. Geochem. Explor.* 67: 407–415.
- BORGNA L., DI LELLA L.A., NANNONI F., PISANI A., PIZZETTI E., PROTANO G., RICCOBONO F., ROSSI S. 2009. The high contents of lead in soils of northern Kosovo. *J. Geochem. Explor.* 101: 137–146.
- DAS A.K., CHAKRABORTY R., CERVERA M.L., De LA GUARDIA M. 2006. Determination of thallium in biological samples. *Anal. Bioanal. Chem.* 385: 665–670.
- DMOWSKI K, KOZAKIEWICZ A, KOZAKIEWICZ M. 1998: Small mammal populations and community under conditions of extremely high thallium contamination in the environment. *Ecotox. Environ. Safety* 41: 2–7.
- ENGSTRÖM E., STENBERG A., SENIOUKH S., EDELBRO R., BAXTER D.C., RODUSHKIN I. 2004. Multi-elemental characterization of soft biological tissues by inductively coupled plasma-sector field mass spectrometry. *Analytica Chimica Acta* 521 123–135.
- EPA Maximum Contaminant Level <http://www.epa.gov/safewater/contaminants/basicinformation/-thallium.html>
- GALVÁN-ARZATE S., SANTAMARÍA A. 1998. Thallium toxicity. *Toxicology Letters* 99: 1–13.
- HEIM M., WAPPELHORST O., MARKERT B. 2002. Thallium in terrestrial environments- occurrence and effects. *Ecotoxicology* 11: 369–377.
- HRŠAK J., ŠKRBEČ A., BALAGOVIĆ I., ŠEGA K. 2003. Thallium Content in Zagreb Air. *Bull. Environ. Contam. Toxicol.* 71:131–134.

- JACOBSON A.R., MCBRIDE M.B., BAVEYE P., STEENHUIS T.S. 2005. Environmental factors determining the trace level sorption of silver and thallium to soils. *Sci. Total Environ.* 345: 191–205.
- JAKUBOWSKA M., PASIECZNA A., ZEMBRZUSKI W., ŚWIT Z., ŁUKASZEWSKI Z. 2007. Thallium in fractions of soil formed on floodplain terraces *Chemosphere* 66: 611–618.
- KABATA-PENDIAS A., PENDIAS H. 2001. *Trace Elements in Soil and Plants*, third ed. CRC Press, Boca Raton, FL: 356.
- KAZANTZIS G. 2000. Thallium in the environment and health effects. *Environmental Geochemistry and Health* 22: 275–280.
- KRASNODEBSKA-OSTRĘGA B., DMOWSKI K., STRYJEWSKA E., GOLIMOWSKI J. 2005. Determination of thallium and other elements (As, Cd, Cu, Mn, Pb, Se, Sb, and Zn) in water and sediment samples from the vicinity of the zinc-lead smelter in Poland. *J. Soils & Sediments* 5 (2): 71–73.
- LEONARD A, GERBER GB. 1997. Mutagenicity, carcinogenicity and teratogenicity of thallium compounds. *Mutat. Res.* 387: 47–53.
- MARTIN H.W., KAPLAN D.I. 1998. Temporal changes in cadmium, thallium and vanadium mobility in soil and phytoavailability under field conditions. *Water, Air, and Soil Pollution*, 101: 399–410.
- MOCHIZUKI M., MORI M., AKINAGA M., YUGAMI K., OYA CH., HONDO R., UEDA F. 2005. Thallium Contamination in Wild Ducks in Japan. *Journal of Wildlife Diseases* 41(3): 664–668.
- MOORE D., HOUSE I., DIXON A. 1993. Thallium poisoning. Diagnosis may be elusive but alopecia is the clue. *BMJ* 306: 1527–1529.
- PAVLICKOVA J., ZBIRAL J., SMATANOVA M., HOUSEROVA P., CIMAROVA E., HAVLIKOVA S., KUBAN V. 2005. Uptake of thallium from artificially and naturally contaminated soils into rape (*Brassica napus* L.). *J. Agric. Food chem.* 53 (8): 2867–2871.
- PETER A.L.J., VIRARAGHAVAN T. 2005. Thallium: a review of public health and environmental concerns. *Environment International* 31: 493–501.
- QI W., CHEN Y., CAO J. 1992. Indium and thallium background content in soils in China. *Int. J. Environ. Stud.* 40(4): 311–315.
- RAO C.R.M., RUIZ-CHANCHO M.J., SAHUQUILLO A., LÓPEZ-SÁNCHEZ J.F. 2008. Assessment of extractants for the determination of thallium in an accidentally polluted soil. *Bull. Environ. Contam. Toxicol.* 81: 334–338.
- RUSYNIAK D.E., BRENT FURBEE R., KIRK M.A. 2002. Thallium and Arsenic poisoning in a small midwestern town. *Annals of Emergency Medicine* 39; 3:307–311.
- RÜHLING Å., TYLER G. 2004. Changes in the atmospheric deposition of minor and rare elements between 1975 and 2000 in south Sweden, as measured by moss analysis. *Environmental Pollution* 131: 417–423.
- SABBIONI E., MINOIA C., RONCHI A., HANSEN B.G., PIETRA R., BALDUCCI C. 1994. Trace element reference values in tissues from inhabitants of the European Union.

- VIII Thallium in the Italian population. *Sci. Total Environ.* 158: 227–236.
- TREMEL A., MASSON P., STERCKEMAN T., BAIZE D., MENCH M. 1997a. Thallium in french agrosystems: I. Thallium contents in arable soils. *Environmetal Pollution* 95 (3): 293–302.
- TREMEL A., MASSON P., GARRAUD H., DONARD O.F.X., BAIZE D., MENCH M. 1997b. Thallium in french agrosystems: II. Concentration of thallium in field-grown rape and some other plant species. *Environmetal Pollution* 97 (1–2): 161–168.
- TYLER G. 2004a. Ionic charge, radius, and potential control rootysoil concentration ratios of fifty cationic elements in the organic horizon of a beech (*Fagus sylvatica*) forest podzol. *Science of the Total Environment* 329: 231–239.
- TYLER G. 2004b. Vertical distribution of major, minor, and rare elements in a Haplic Podzol. *Geoderma* 119: 277–290.
- VANĚK A., CHRASTNÝ V., MIHALJEVIČ M., DRAHOTA P., GRYGAR T., KOMÁREK M. 2009. Lithogenic thallium behavior in soils with different land use. *Journal of Geochemical Exploration* 102: 7–12.
- WANG Q., HUANG X., LIU L. 2007. Analysis of nine cases of acute thallium poisoning. *Journal of Huazhong University of Science and Technology* 27 (2): 213–216.
- WIERZBICKA, M., SZAREK-LUKASZEWSKA, G., GRODZINSKA, K. 2004. Highly toxic thallium in plants from the vicinity of Olkusz (Poland). *Ecotoxicol. Environ. Safety* 59: 84–88.
- WINTER M. J. 2007. WebElements the periodic table on the WWW, <http://www.webelements.com/>
- XIAO T., GUHA J., BOYLE D., LIU C.Q., CHEN J. 2004a. Environmental concerns related to high thallium levels in soils and thallium uptake by plants in southwest Guizhou. China. *Sci. Total Environ.* 318: 223–244.
- XIAO T., GUHA J., BOYLE D., LIU C.Q., ZHENG B., WILSON G.C., ROULEAU A., CHEN J. 2004b. Naturally occurring thallium: a hidden geoenvironmental health hazard? *Environment International* 30: 501–507.
- YANG CH., CHEN Y., PENG P., LI CH., CHANG X., XIE CH. 2005. Distribution of natural and anthropogenic thallium in the soils in an industrial pyrite slag disposing area. *Sci. Total Environ.* 341: 159–172.